





UNIVERSITY OF CALIFORNIA
MUSEUM OF VERTERRATE ZOOLOGY







Boyers, L.M. Freeno and Kern Counties, Calif. 1932



Boyers, L.M.

Fresno and Kern Counties:

Fowler

1/2 mi. E Miramonte (F. Co.)


18 mi. E Bakersfield (K.Co.)

Bodfish (K.Co.)

Waltham Creek, 41/2 mi. SE (Riest Cally (F.Co.)

Dec. 15-29, 1932

Specimen nos. 291-361




(291) December 15, 1932

Robert T. On and I left
Berkeley at 9:30 A.M. this morning
and went out the Tunnel Rd. the
Dublin, then through Tiermore ralley
over alternant Paks and through
Tracy, Manteca, and down the San
Joaquin Valley to Fowler by Modesto
and Fresno. We are stopping thright
on the southing limits of fowler.


Miramonte, 3500ft, Fresno Co., Calif. - 1/2 mi E- at Dan Crossing December 16, 1932

This morning we left Fowler after getting a few supplies and went to the outstants of the dy, where we turned off east to go thru Orange Cove them to Miramonte (3500ft.) and then I mi further to Deer Crossing where we have taken a small cobin. After arriving in camp about 11:30 A.M. and getting lunch, I started out worth of camp to explore and set out some mouse traps later.


The most common tree in the




M. Boyers
1932 1/2 wie F., Miramonte, 3500 ft., Fresuro Calif. December 16, 1932 camp is the yellow sine (Cirus pouderous) with scatterings of Buck-eye, Madrow, Manganeta, small leafed live Oak Quereus dumora, Q, wistgener, a Hahoforicarpus albus, Incense adas (1) and supposedly more 1/ N.E, also White Fir supposedly cupeatus 1/2 N. E of comp. also Phannus crocea, Ceanothus, about a / mi N. of carry, on my trip This afternoon, I saw, for a fraction of a recond, a Weasel running along a fallen log, partly open to view, and their disappear in a thick brush patch. Hylocichla guttata's ar very numbrous, ils fact almost the most common brid present, will Passerella ileaca a close recond, then the bids seen or heard were: Sialia sp? (2) head Merula (20-30) saw Sycreus M. (4015) saw Pipilo fuscus (6-8) saw, heard, collected two Pipilo erythrophthalums (3-4) heard ashelocoma californica (2-3) saw Buts virgineanus (2) and, head Tophortyx californies (3-4) saw Chamara f. 1/1) heard I ryobates villosus (2) saw




L.M. Boyers 1/2 mi. E., Misamonte 3500 ft, Fresul Co., December 16, 1932 Isaw several wood rate nests, but nothing that looked very fresh, Deer tracks, brush robbit tracks, for tracks, and coyote tracks were exceedingly common. On the last gart of the trips I set out 35 mouse traps. The ground is covered in the protected spots with from two to four wiches of snow, that supposedly fell thee or four days December 17, 1932 Up at six thirty, buakful, and out on the traps line, but at ramed during the night and many were spring and the others had reothing in them. a large part of the snow was welled by the rain. Ho new birds were seen or heard except a Colaptes calendula, one of which I shot. after hunting variety for brush rabbits for about two hopes I returned to camp and shot a Pipilo fuecus on the way. I put up one bird and then got lunch, after which I spent until to clock



1/2 mi. E. Misamonte, 3500 ft., Fresno Ca, 61 M. Boyers 1932 December 17, 1932 putting up another Pypolo fuscus and examining and preserving infected parts of Orrofour mice (Peromyseus calf, and 3 Peromysons boylis). Blood smears were made of both of my Orjailos and Orrs 4 mice as well as a Tasserella iliaca that Our putup. One Peromyeeus boylii (871 8-On) had an abovess (1.5 m. m dia) on the right central lobe of the liver and the bidneys appeared to be finely mothed on etched. The intestine appeared to be normal. Wt 24,59 Another Peromyseus boyli (8738-On) had about 15 minute abscesses around the thin edges of the lobes of the lever. The bidneys and intestins appeared to be normal! (No intestinal severas were made of either of the above 2 specimens Wt. 27,29. When retreening the Sipile Justus I shot this morning, I came across the whole body (skinned) of at best as a sich up (). In talling with the man in charge of the camp here at offer Crossing, (Mr. Willes) I learned that a wearef




1/2 mi. E. Misamonte, 3500 ft., Fremo Co., 62 M. Boyeve 1932 Calif. December 17, 1932 had been taken about a week ago, but was thrown away, without even taking this skin off. The conversation with the people here, espeadly those that trap during the winter, we have found out that locally the forblowing for animals are taken; Mountain leon-Bob cat-Coyote (Vally) They Fox (common) Red Fox (raver) Stryed Skunk (common) Sported . " Weasel-Brush Rabbet (not common) This evening from 4 \$ 5730 when it was dark, I hunted up a creek bottom that was bordered on one mile by a meadow with farry high grass, partly under snow and shot another Pipilo fuscus and saw and heard Hairy wood sectors, Cobins, Latif. Jays, and Hermit Itausker. a live of 35 trayes was set along and bordering on the creek, some in the snow are some not.




December 18, 1932


Up at 5:50 A.M. and went rounds of traps, getting a Berongseus boylis and a Veromyseus trues, then went hunting brush Babbits, saw two, but got weether. They were on the shaded, west facing stope, pussing under low, thick, sprawledout patche of Madrone and in both cases were in the open only a faction of a record as they crossed a list trail. Voison oak and Manganeta as well as Buckeye formed almost impassable thickets, growing down to within twelve to eighteen inches of the ground. While looking for rabbits I saw a Grey Squiriel come down a large Black Bake tree and go off the the bush. We skinned until about 2:30 P.M. and then went in the truck down (west the road to a place / mi W. Musmonto (direct distance) and set out about 20 traps apace around some promising holes that anay be Veroquathus holes. They were from an inch und a quarter to two inches wide and were left open as well as slunted gently into the ground. No drappings were found.




1. M. Boyers /2 mi. E. Minamonts, 3500 ft., Fresul Co.,4
1932 December 18, 1932 after that we went to a place 3 mi. W. Meromonte and set out traps for Gerompaus Calif in moss-covered, boilder-strewn, Madrone and Muszamito covered slopes. Wood rat sets were also made in some of the frequent wests. a Western Gratcather was seen here as well as a Winter Wren (1mi W. Miramonte) to add to the bid list. On also reported Evening Grosbeales and Gewich Wrens. We heard Wentets and Olaintiturouses, but so far haven It been able to secund any. Ho brush robbets were seen att either of these Bratisis West of Mirawoute. also those I saw this morning were moving down hill at between 8:00 and 8:30. The sur was just reached, the crest of the hill and they were coming down from the ridge proper toward the still shaded canyon. Heither specimen 296 on 297 had infected livers on beidneys on intertures by external exacus in a trois and Oa's opecimens 877, 878, 4879 were not in fected in these same areas. Glood smears were made. The weather was warmer today



1/2 min. F. Miramonte, 3500 ft., Frems Co., 65 1. M. Boyers 1932 December 18, 1932 and the snow is going very fast, only the most protected spots till being covered. also at the location / mi. W. of Miramonte where we set the trops for Verognathus, I heard and saw one Cutamas in a small pile of large welks on a shaded North facing slope, just offthe road. Efforts December 19, 1932 Up and dressed by 6:00 A.M. this monning Tot in the truck and went W. to och traps I-wi. W. of Miamonts, where On got one Peromyseus calif and I got one Peroquothus calif. that love out the theory about the holes being Ceroquother holes. at this location we saw and heard a Writer When and White-crowned Spanows. at the second location, 3 mi W. of Misamente, 2900 ft, Ors got one Brognathur calif on a shaded brush covered hillride at the foot of a black oak with moss covered rocks around.




. M. Boyers /2 min E. Miramonte 35-00 ft., Fresno Co., December 19 1932 The trap was set right ment to the beginnings of a Geotoma pel. This one specimen was the only one taken at this location, Our and I separated and hunted for two hours, Or getting a Plain Titmouse, Spotted Towher, Brown Towher, Wren-tit, and a Juneo, also one Brush Rabbit. Woth of us saw a flock of about 120 Band-tailed Pigeons Plying Fast overhead. I seewed a Downy Woodprelser in the top of a Black-oak along a creek bottom that was lined with Calif. Lawel and Willows . Shortly afterward ? secured a Bewick When and a When-tit in a taugh of falley Buck-eye and manguneta. The win-tit was occompamed by a second, but it flow so close, while I was wrapping the first up that I couldn't shoot it without flowing it to pieces. I wan a brang Spiritel in the top of a tree, but due to the denseness of the Soush setween couldn't kill it with our shot, and I didn't get a second. I found a finall stram-cut gully man dry, that had linge




M. Boyers 1/2 mi. E. Miramonte, 3500 ft., Fresno Co.,

Calif.


December 19, 1932 boulders in the stream bed and projecting from the sides and covered with mos and snow. Got holes were worn in the rock in the stream bed and these holes were about 2/2 ft in dameter and fire feet deep, Brown Towhees, Spotted Towhees, Ruby crowned Kringlets, Plain titmouses occared here. Tater going back to the cart I shot another When-tit and a second that was with it. In fact they seem to always be found in pairs. at four P.M. we went back and Our hunted brush rabbits while I set traps for Pergualtius and Dipodomys on open grass covered hillside, bordered by Manganeta and Madaone. I saw one Cotton tail rabbit and six may Band-tailed Digkons. It rained some this afternoon and is fairly clear and warmer to might it completely, I want to record the fact that the frast thinging we heard on arwing 3 mi W. of Minamont, 2900 ft., was two John. Threshers singing. Heither of us was able to get a specumen er even hear or seg




1/2 mi. E. Meramonto, 3500ft, Fresno Con Calif. December 19, 1932 another Thrasker. The location when we set the trajes today was typically Upper Souveret, with open hillordes bordered by bush patches of Buckeys, Maryaneta, and Madrone, as well as seatherings of Oak (Blacks and a small leafed tak-). along the creek there Willows, Wild Black berry and Lawel in the shaded, wetlest parts. The bush is more open here than at the other locations where us have been hunting. December 20, 1932 Ely at 6:20 and had breakfast then went 3 mi Woof Miramoute, 2900 ft, to our trapping location. a heavy fort covered everything and it was extremely cold. No Berognathus or Sypodowny wey cought, but one Peromysus maniculatus was in our of my mouse tropes We spent they down hunting birds, nuther of no seeing any rathets. I secured a Fox spanow in the manganeta funch on an east facing slope as it scratched in the dut and leaves on the grown Ment Is shot a Hairy Wood secker in a



1/2 mi E. Misamonte, 3500 ft., Fresus Ca, M. Boyers December 20, 1932 Black Oak on the creek bank. The first shot, a no. 6. shell, but the woodpecker in the upper back and base of the skull and penetrating it, yet the bird lung backwards, hoad back and down, by its feet. Two 10.12 shells had no effect, not way soutrating the body. Finally after about fut minutes as I was about to give up the bird fell, still alis and had to be helled by hand They as certainly tenacions to life. On skinning this was explained another icily by the fact that the nech is lasily wiset since it is so their, the head is hard to penetrate due to a thick should, the back is formed of very heavy vitebrae, and the breast uniscles and sternum extend almost to the auns, leaving only a small unprotected area of viscera. Then too, the bread mussely are abnormally large for the size of the bud. great I secured a Plain Titmona out of a small leafed daloa City Jay as it flew over head with a linge acom in its mouth, and



1/2 mi. E. Miramonts, 3500 ft., Franco Co., 70 Calif. December 20, 1932 . M. Boyers last a Juneo as it flew out of a bush on an open bill sidt. On the way back to camp not noticed that the weather was actually very warm and surney, and as wel drove along I sow a Calf. Ground Squirel wear it, hole at the side of the road and I secured et. about a mile further on we saw a second one, but couldn't get it as it ran down its hale with a Ducke eye ball in its worth. December 2/1932 It raised and bailed cluring the night and started snowing about siyo'clock and by ten o'clock we had from the to four inches over everything. We broke camp and after trying vivily to get chains on over sized ties, we pulled out and had no difficulty ingetting to Redly. On the way down a Poison Out "note and Tewis Twood prekers and later Melering Birds in numbers. It raised off and on all day,



18 mi. L. Balsersfield, entrance to Seguoia M. Boyers 1932 Mat'l Forest, Hern Co, Calif.

December 21, 1932

We finally reached Bakersfield Tra Fulare (where we had bunch and

stocked groceries). In Bakersfill we went for weil and got a few

supplies then beaded out the

Kenwille road and are campel Tought about a fundred yards south

of the Kennier at the mouth of the Kern Canyon, I set out about

25-30 traps in the flat ground

and down a erets wash. It ramed


hard a couple of tunes in the process of getting a meal and


now has stopped again. It is not

arry cold.


December 22, 1932

While out getting traps and during breakfast, the finds over and heard were: Great Blue Jerou, flying east up the Kerry; Rock Tekens, out on the flat and on the creeks banks; White-crowned Sparrows, in the brush lined dry washes; Timet, ou the river bank; Puby-crowned Kinglet, in cottonwoods along river edge, Horned thats, out on the flat weed covered or open land; and finally a Red-tailed Hawle as we left and entered







Bodfish, 2600 ft., Kem Co., Calif. .M. Boyers 1932 December 22, 1932 Frank Poss spends most of his Line bunting cats (Bob and Lion) with a gack of Thounds. He trains them himself, taling two to three years to produce a good how deg. His method of trowing is to trap a bob-cat, gather the young dogs and one experienced dog, released the cat out of sight of the dogs, let them see and swell the fresh cut tracks and follow it under the guidand of the old dog. after has 3 hunts they pich up a lot of Expensive at it and ban rapidly. He has gotten from 90 to 125 for one of his trained him dogs. F. Ross till finds liver in this general region and has all the people notify line on a cats first appearance. J. Bruce has left this region to him for a number of years. Dear are still to be found, but both Ross brothers class them as hamles, saying they will not bill cattle, but will eat fallen game of any brut, trampling out all other tracks and making it look like bear works. J. Rosse asserts that would the bar in this region actually hibernote Thru a writer, but next up temporably during




Bodfish, 2600ft, Kern Co, Calif. L. M. Boyers 1932 December 22, 1932 storms, but can be found out otherwise. He says their beds are basket shaped, about six feet long and four fut wide and are made entirely of price needles. Leces are dropped in a cerèle about the Melst in a heavy ning. It was of interest to note that the only weapon! Ross carries when lion hunting or talsing a party out ofter bear, is a .38 Police positive revolver and high power ammunition. Says new high power ammunition is better than 45 shelf for shocking power. always shoots hour from below, when they are treed, shooting for med-chest and have bulket range upward then neck and basa skul region. F. Ross and H. Ross both thirls that predatory game control by paid gov't trappers is bad in more than one way. I. Hoss said he followed up a creeks where a game control agent has been trapping and found 19 striped Showshe billed and thrown away, He has seen endered of xinsilar throwing away of countities grey fox. Impast years used by get samue, gren for in numbers and a flw-coyote, now are getting






M. Boyers Bodfish, 2600 ft, 18em Co, Caly, 76 December 22, 1932 recalled W. P. Taylor and firstay there, collecting in 1911. appreciated belg girly them by M.V.S. on several questions so which they had asked about. F. Poss recalled seeing a large "blue bullet hawke" with a wing spread of about four feet or word attacking a red-tailed hawks a mumber of years ago in a tree in front of their colvin (Evidently a Joshawls) Both appeared sincers and rebable in every respect and their anouted ge of aminals was unlimited as for as local game was concerned. H. Ross mentioned "grunting med" in the calrie not lenowing whether it was caused by lenaving or by a word effort. Recalled three eagles attacking a dol and attempting to tout it until driver off. Fefstrough, against protecting eagles in this region saying they ful certain that eagles get maky young favus as evedenced by funding eagles on freshly billed faire bodies. Both brothers had seen white-tailed jack robbets years ago up is the higher





mountains. HowRoss called them Southernsierra Hores and benew of four members of ette robbit series, Pilsa, Brush robbit, Cottontails, + gads-brobbets, having seen them all during the time of trapping. Realled watching tissas stack hay and of examing a live one once. Both barely able to she out a living as fur is bery low now. F. Rosssent in 3 coyote skins and got a check for 65%, while Hiloss sent in & prime coyote skins and got only 509 agrice for them. He should me the fur companies bill on it. He has only been getting about 11. for his grey for pelts and strangely enough from 1.50 to 2.00 for Bob cut which is usually forver than the

The Ross has kept several hors
cubs and mentioned black, curling
shoulder stripes running down behind
the front legs when the cubs are young.
I. Poss spoke of seeing many condors
in past years, ging a good description
of the white inderwing patch and the
yellow-leathery necks, waked of pathers.
He said that in 3 hrs. he could still take




Bodfish, 2600ft, Kern Co, Calif. M. Boyers 1932 December 22, 1932 us with condor country on an all road, that they frequently travelled that lay be tween Balsersfull and J. J. Coss also stated that he had been told authoritatively by gov't trappers and possoners that they had orders to report only 2 or 3 grey for taken each year und no more to give a right appearance" to the record, since no foxes caught would look que. Consequently mumerous grey for we chucked, and only the "proper no "reported. Just another black eye for control measures. after getting the test up and the equipment organized Orr and. I set out traps. I set a line up over a slope curered loosely with Juniper, Yucra whipplei, Jurcus duriosa, and Ceanottino a scattering of dead grasses. I primped a Cotton tail out of a Juniper clump and got it with two shots. It disappeared under a second Juniper clump 50 yds frither up the







Bodfish, 260 oft, Pern Co., Calif. 80 M. Boyers 1932 December 23, 1932 of a Barn Out hung on a barbed wire fence, Titmouses, Tubycrowned Kinglets, Mexican Blubrids, timeto, and Juneos. Shot two Juneos from a feeding flock on an your weed covered hillseide of about 150 to 200 brids. Oar out up oul of the Juneos, I also secured one Bluebird. It ramed, hailed, and snowed interchangebly all day and finally let go with a good wind making life in a tent anything but pleasant. We decided to pull out in the morning as north snow-covered, brushy slope yulded not even a snigle rabbit track to Orro best efforts. Pablits higher up if any are present and this is nin passable for us at present-December 24, 1932 Feft at about noon after talking again with the Rose brothers and headed back to Bakersfield. We stocked up and left from there for Julan where




Waltham Cr., 4/2 mi. S. E. Priest Valley, 81 M. Boyers 1850 ft. Fresus Co., Calif. December 24, 1932 we lunched and laid on more supplies. Went morth to mean Soshen them turned west the Coolinga, then toward Priest Valley and finally at dark made arrangements with T. B. Green to stay on his ranch in and old store building on his ranch. It was doo dark to set out any trapa so we planned to get out early after rabbits tomorrow. December 25, 1932 In working over upland plateau, oak covered ground got one Cottontail and several brids, sitta c aculeata, 2 Zonotrichia L. gambeli, 2 Baeolophus inormatus, / Junes oregonus, and one astragalians p. herperophilus Inever saw so many Tawence and Green backed Goldfinches or heard such a drive in my life. They were present by the hundreds, it being common to see from 20 - 30 in one the top. according to T. B. Sheen and his nephew A. A. Bonds this is the usual thing. I shot one of the funches and discovered it was




Wattham Gr. Hz mi S. E. Briest Valley 82 C.M. Boyers 1850 ft, Freams Co, Calif. Deember 25, 1932 a Green backed female. They seemed to be mixed in with the Faurence Goldfriches. In the late afternoon we both set out traps after hunting for rabbits. I saw one Cottontail, outlined against the sky on an open slope 75 yds. away when it had become almost totally dark. I set out about 35 trapes over the Oak covered high-up-flats and set some around the Peroquetteus holes that are so numerous on the semi-open oak spotted slopes. Set out total trap him when it was sitch black, by flashight and a heavy mist came in obscuring objects over 20 yds away. Fot hung up on gulch brunks, bordered by a second and didn't find my way out for a good helf hour. Saw out live Dipodomys on way down tried to get it alive but couldn't. Our did succeed in gettering a his one by setting a couple of traps on either side of a bush and driving the Dipodomys mits one of them. The was cought by one fort.




Waltham Or, H/2 mi. 5. E. Priest Valley, 83 1850 ft., Fresus Co, Calif. M. Boyers December 26, 1932 Went over trap his after seeing no rabbits and only secured 2 Peromyseus truei and 2 Peroqualtus Californiens. Have seen quite a few brids here: Valley quail, Mourning Dow, Calif acomstoring Wood sicker, Trimet, Mexican Bluebird, Laurence and Green-buckey Told friches, Hermit Thrush, Rubycrowned Kinglet, Claint tuouse, Bushtit, Robin, Calif. Jay, West tailed Hank, Cooper Hawle, Sparrow Hawle, Slender-billed White-breasted Thuthatch, Junco, Samle to hito-crowned Sysavan; Solden-crowned. Sparrow, american Crow Varied Thrush, Calif. Known Tawher, Spotted Towher, Red-shafted Flicker, Larlo Sparrow, Tog gerhead skrike, Calif. Thrasher, Vigors Bewick Wren, Wren - tit, Say Phoebe, Fox Sparrow, Western Grateles and american light (34 species) Spent major part oflay doing examin ations and blood smures of accumulated material. Strangely enough & haven't discovered another abscersed lives series the one obtained from the catch at our prost camp a/2




Waltham Cr., 4/2 mi. 9. E. Priest Valley, 84 M. Boyers December 26, 1932 mi I. Miramonte, 3500 ft. Fresno Co. about 4 o'cleck or shortly after, On and Dwent up to the oak flat that is covered in places with adenoitoma fasiculatum, Ereogonum, Ceanothus cuneatus, Quercus dumosa and Cercocarpens Betuloides, also alluin, Browns There we separated and set out traps in a large Ripodomya colony that covered the top of one whole butt, The burrows slant into the ground at an angle less than at the entrance, usually strong out into a small slanting mound in front of hale. Top view Side view The openings may be in a worn open place or by a dead fallen limb or in dead grass or near small bushes, and are commented up for distances of 15 ft. on man




Waltham Cr., 4/2 mi. S.E. Priest Valley 85 L. M. Boyers 1932 1850 ft., Franco Co, Calif. December 26, 1932 by paths worn then the dead grasses and welds. These may cross and recoss, but always take the shortest distance between two holes, I set out 50 traps on the colony flat and 18 more gart way down the trail to camp. White putting out traps 9 saw a Black-tailed Jack Rubbit and be headed from the open who the close thick growing adenostoma fasiculation In fact both cotton-tails and Jacks seem to prefer this adenostones since all of both I have seen have either come out of or gove into it. December 27, 1932 Some of the other plants and bushes seen here that are fairly common and Symphonicarpus sp?, Solanuin Kanti. var. violermedium, Salvia, agrostis, and Erio dyctyon crassifolimis Took in tropo this A.M. and paw 2 more Jack . Rabbits . and again when both headed a cross an open flat over 100 ydo long and plunged mit an adenostoma thickets




L.M. 13 ogers, Waltham Cry 4/2 mi. S. E. Prest Valley 86 1850 H, Freamo Co., Calif. December 27, 1932 Got 3 Dipodomys on the colony location and one more on the down trail. also secured 5 Bromyseus trues from among the Dipodomp holes at the colony est nations likewise 9 got 2 Peromyacus manuculatus on the down trail. Spent day skinning. according to 1, B. Green and other local inhabitants this area was burned over several truts with in the last find years, and the farmers in general consider brush rabbits and cottontails, and Jacks to a leaver degree, as downright pests, saying in one case a fill mear a brush his had an 8 ft. strip cleaned off during the anghts, also on what appears to be good abouration, the was à disease :, at least a market decrease ni the rabbit population in the young of 1930. December 28, 1932 Heavy frost this morning; went bird hunting and saw few birds surce a heavy wind carrie up and they all kept close to cover.




M. Boyers Waltham Cr., 4/2 mi. S. E. Priest Valley 1850 ft, Freano Co., Calif. December 28, 1932 Gat a Blue bird, a Olain Tit mouse, and a Tambel white Crowned Sparrow. Did more blood smeans and then gut up 2 Ononyocus trusi and 3 Perongeous marriculatus that On secured. Bird ortuation books pretty bad as far as no. of specimins oftamed. T.B. Green has a mud mind about ten feet long and 3 feet wide that oozed out on anopen hillorde. It crustsover in dry weather, but is always soft! underweath. The mind like clay is an olive brown color and / contains no grit. Only washing ones hands in water with a little of this clay for some leaves the hands faultlessly clean and puits, even if they were greasy, December 29, 1932 Went bird hunting and got 1 Larrino budovicianio, Bacolophus siromatus, I aphilocoma calif., and 2 Pypilo fuscus. Oldent around gopher traps with arr jegetting some, and orr also getting a clipmuk



M. Boyers Waltham Cr. 4/2 mi. 5 . F. Priest Valley 88 1850 ft, From Co., Calif. December 29, 1932 by shooting it off of a fallen logina dry wash. I also got Two more Bacolophus inornatus. While eating breakfast on the gorch 9 shot a Cargodacus mexicanus find another Lanus budovicianus from the top of a valley oak a measured distance of 55 ydo with a 410 shotgun, and a no. 8. shell, Saw one brush rabbit run from under a Photoma arbitifolia down but them loose brush, and wasn't able to secure it. Brush robbits and cottontails both inhabit semi-open brush country. here and also the Jacks sto a lesser degree. A. H. Bondo brought in a Pyperlo fraces and a Bysilo maculation This country is covered in places with shell deposits and T. B. Shen showed no rib fragments and vertebras over 6 minches long and Hor 5 muches this, petrified, that had been picked up on his raich.



M. Boyers Waltham Cr., 4/2 mi S.E. Priest Vally 89 1850 ft., Freamo Co., Calif. December 30,1932 Got out at 1:15 P.M. and went over to Onest Valley, seeing large flocks of meadow larks and up Mustang Grade to San Tucio, there to Salvias and Jan Jose. From there to Haywardo, Oakland, and Berkeley by 8:30 P.M. Saw 2013 Paraie Falcous ditien coming down the west side of Mistang Grade nito San Sucas.




Wt. 68.9

Wt. 98.59 Wt. 19.49

4 mi. E. Miramonte, 3500 ft., Fresuo Co.	Calif.
December 16, 1932	
29/07- Ripilo fuscus ot	Wt.56.39
2929 " 4	W1.51.3g
December 17, 1932.	
2938 Regulus calendula d	w. 1. 6.49
2949? Pipilo fuscus 9?	WH. 16.39
2959 " " 9	WH.55.19
. December 18, 1932	
2960 Peromyseus boyli d 173-83-22.5-19	Wt. 24.9
2978 " truei of 156-83-22.5-21.5	Wt. 16.6g.
Lui, W. Miramonte, 3200 ft, Fress Co, December 19, 1932	, Calif.
2989 Perognathus calif. 208-121-26-11 200 Dyobates pubescens	W1.2629
300 & Chamaea faciata	Wt. 14.89
3 mi. W. Miramorte, 2900 ft, Fresno Co., Ca	lif.
December 19, 1932	
30/0 Chamala faciata	Wt.16.69
3020 " December 20, 1932 3038 Basolophus inormatus	W. 15.59
3636 Basolophus mornatus	WH. 15.29

304 & Dryobate villosus 305 & aphelocoma california

306 & Juneo oregonus



Becember 200 ft, Franco Co, Calif. December 20, 1932

309 & Peromysous maniculatus 172-84-20-19 W.17.49 308 & Citellus beechyi, 413-177-60-27 W.639.29

18 min E. Bubersfield, mean entrance to Sequeria Mat'l Forest, King Co., Calif. December 22, 1932

3099 Peromyseus maniculatus 162-17-185-17 W. 18.69
31001 " 148-67-19-15.5 W. 15.49
31107 " 155.5-70-21.5-17.5 W. 18.69
31201 " 149-70-20.5-18 W. 17.69
31301 " 142-65-21-16 W. 12.89
3149 (mursing) " 160-70-20-16.5 W. 18.39

Bodfish, 2600 ft., Kensler, Calif.

December 23, 1932

315 & Peromyseus truei 197-113-24.5-24.5 Wt. 25.29

3/6 & Peromyseus boylii 172-90-20,5-19 Wt. 20.89

3/1 of Peromyseus maniculatus 149-68-20-16 Wt. 18:29
3/8 of Sialia medicana occi. Wt. 29:89

3199 Juneo oregonus

Waltham Cr. Hui S.E. Priest Valley, 1850 St. Fresno Co. Calif.

1850 ft, Fresur Co. Calif. Delember 25, 1932/ C= 96 320 & Sylvallagus anduboni 400-63-94-31


32/4 Sitta c. aculeata

32201 Zonotrichialgambelii

W. 158.9

W.17.69

Wt. 18.79 Wt. 26.49



Waltham G., 4/2 mi St. Prest Valley, 1850fl, Tresno G., Calif. December 25,1932


323 of Lonotrichia l. gamble	WH. 25.99
324 & Bacolophus inornatus	WH. 15.19
3259"	WH. 15.79.
326 of Junco oregonne	Wt. 18.29
3274 astragalisms facourer hesperophilus	

December 26, 1932

328 of Perompen	s trues	204-107-24-25	WH. 29.59
32907 "	"?	176-82-23-21,5	Wt. 21.79
330 & Perognathus		205-113-26-115	Wt.22.69
33/7 "	· ·	210-115-26-11.5	W1.25.89

December 27, 1932

332 8	Dipodomy	heerm	ani 300-195-42,5-17	Wt. 76.89
3338	"	"	295-175-43.5-16	Wt. 87.39
3349	4			Wt. 67.9
335 8	"	Manicula	300-184-46-17	Wt. 88.49
336 8 7	Peromyseux	M. Call Market Call Ada, M. Cal	154-68-20.5-16.5	W# 2/.9g
3379	11	tout the same of t	178-91-23, -23,5	Wt. 26.49
338 8	"/	46	202-110-23.5-24	W. 29.19
3398	′,	4.	09-(75)-24-23,5	Wt. 24.49
340 8	',	q :	167-80-22,5-21	Wt. 22.39
34/8 Pe	romyseus ma	minhatus	144-59-20-16	Wt.15.6g
3420		"	124-55-19-16	Wt.13.29



3 Waltham C., 4/2 mi. 5. E. Prest Valley 1850ft, Fresuo Co, Calif. December 28, 1932

343 & Zon	otrichia l. gambeli.	Wt. 22.19
	lophus inornatus	Wt. 16.69
	wens trues Coll. R.T. On 190-94-22.5-22	W1.24.69
,	" " 173-83-23.5-22.5	W1.20.69
3479	mamentatus " " 159-66-21-16	Wt.18.89
348 2 "	11 11 156-68-20-14	Wt. 18.39
34981 "	" " " 152-65-20.5-16.5	W. 20.9

December 29, 1932

3509 Tanius ludoviciams	W/t. 508g
35/9 "	Wt.55,29
352 8? Bacolophus inormatus	Vt. 17.39.
3538 "	Wt. 18,59
354 9 "	Wt. 16,29
355 q aphelocoma calif.	W.83.29
356 & Caryodacus mej camis	Wt.18.59
3570 Pipilo fuseus	wt. 53. 59
358 9. " "	Wt.49.79
359 8. 11 11 Coll. A.A. Bonds	wt. 58.49
3600 Pipilio maculators " "	Wt. 41.59

361 Wroeyn calif. Picking-December 21- / mi. E. Minawet. 3500 ft., Fresus Co., Colif.

Collecting Stations

1 - 1/2 mi. e. Miramonto, 3500ft, Fresno Co., Calif.

1a-3mi.w. " " " " "

2 - 18 mi. e. Bakersfield, near entrance Sequoia Nat'l For Kern Co., Calif.

3 - Bodfish, 2600 ft., Kern Co, Calif.

4- Waltham Cr., 4/2 mi. s.e. Priest Valley, 1850ft, Fresno Co, Calif.

